

Geometry of weak lensing of CMB polarization

Anthony Challinor^{1,*} and Gayoung Chon^{1,†}

¹*Astrophysics Group, Cavendish Laboratory, Madingley Road, Cambridge CB3 0HE, United Kingdom*

Hu has presented a harmonic-space method for calculating the effects of weak gravitational lensing on the cosmic microwave background (CMB) over the full sky. Computing the lensed power spectra to first order in the deflection power requires one to formulate the lensing displacement beyond the tangent-space approximation. We point out that for CMB polarization this displacement must undergo geometric corrections on the spherical sky to maintain statistical isotropy of the lensed fields. Although not discussed by Hu, these geometric effects are implicit in his analysis. However, there they are hidden by an overly-compact notation that is both unconventional and rather confusing. Here we aim to ameliorate this deficiency by providing a rigorous derivation of the lensed spherical power spectra.

The re-mapping of temperature and polarization anisotropies by weak gravitational lensing can lead to observable distortions in the cosmic microwave background (CMB). In Ref. [1], Hu presented an elegant harmonic-space method for calculating second and higher-order statistics of the lensed CMB temperature and polarization fields beyond the flat-sky approximation. Hu showed that to compute the lensing effects on the CMB accurately it is necessary to use the full-sky power spectrum of the lensing deflection potential. Simple expressions for the lensed power spectra in an all-sky formalism were derived, and their use was advocated for analysis work.

Although we agree with the final results presented in Ref. [1] for the lensed power spectra, we regard several steps in the published derivation as rather confusing. In particular, the re-mapping of the primary polarization on the sphere caused by gravitational lensing is only implicit in Hu's analysis [although it is given explicitly for the flat-sky case in his Eq. (43)]. The point we wish to emphasise here is that on the sphere it is no longer appropriate simply to displace the Stokes parameters (defined on the polar coordinate basis) by the deflection vector; this operation is not covariant and would not maintain statistical isotropy of the lensed fields. This potential confusion is not helped by the overly-compact, and unconventional, derivative notation that Hu has introduced in Ref. [1] without comment. For example, in his Eq. (65) if the derivatives acting on spin-weight 2 objects are interpreted as usual covariant derivatives on the sphere (as suggested by the notation ∇_i), then the lens re-mapping one infers from his Eq. (64) is a simple displacement of Stokes parameters in the polar basis. Here we develop the appropriate re-mapping to describe lensing of the polarization field on the spherical sky in both the tensor and spin-weight formalisms. We then proceed to give a rigorous derivation of the all-sky lensed polarization power spectra that we hope will eliminate any further confusion that may arise from the presentation in Ref. [1].

In the Born approximation, the lensing deflection is

calculated on the unlensed line of sight so the lensing map is a local function of the deflection vector $\alpha^a = \nabla^a \psi$, where $\psi(\hat{\mathbf{n}})$ is the lensing potential (e.g. Ref. [2]). Even for a scalar field, such as the CMB temperature anisotropy $\Theta(\hat{\mathbf{n}})$, there is some ambiguity in how to interpret the deflection vector on the spherical sky. Expressions like $\tilde{\Theta}(\hat{\mathbf{n}}) = \Theta(\hat{\mathbf{n}} + \alpha)$, relating the unlensed temperature $\Theta(\hat{\mathbf{n}})$ to the lensed $\tilde{\Theta}(\hat{\mathbf{n}})$, are only unambiguous in the tangent-space approximation: $\tilde{\Theta}(\hat{\mathbf{n}}) = \Theta(\hat{\mathbf{n}}) + \alpha^a \nabla_a \Theta(\hat{\mathbf{n}})$. However, to calculate the lensed power spectra to first order in the deflection power we need the mapping to second order in α^a . The choice made in Ref. [1], albeit without comment, is to take

$$\tilde{\Theta}(\hat{\mathbf{n}}) = \Theta(\hat{\mathbf{n}}) + \alpha^a \nabla_a \Theta(\hat{\mathbf{n}}) + \frac{1}{2} \alpha^a \alpha^b \nabla_a \nabla_b \Theta(\hat{\mathbf{n}}) + \dots \quad (1)$$

The geometric interpretation of this mapping is that one should displace from $\hat{\mathbf{n}}$ for geodesic distance $|\alpha^a(\hat{\mathbf{n}})|$ along the geodesic through $\hat{\mathbf{n}}$ that has initial unit tangent vector $\hat{\alpha}^a(\hat{\mathbf{n}})$. The mapping in Eq. (1) has the desirable property that it is local in α^a (i.e. the lensed field at $\hat{\mathbf{n}}$ only depends on α^a at that point), consistent with the Born approximation.

For polarization, simply displacing the Stokes parameters according to Eq. (1) is a coordinate-dependent operation and hence unphysical. This would show up in spurious correlations between the multipoles of the lensed polarization field that should vanish for a statistically-isotropic field. The appropriate mapping for CMB polarization involves parallel transporting the linear polarization tensor $\mathcal{P}_{ab}$ (whose components in a specific orthonormal basis are the Stokes parameters) back along the geodesic generated at $\hat{\mathbf{n}}$ by the unit tangent $\hat{\alpha}^a(\hat{\mathbf{n}})$. We can only apply Eq. (1) directly to the Stokes parameters if they are defined on basis vectors parallel and perpendicular to the geodesic at $\hat{\mathbf{n}}$ and its image point.

To establish the appropriate analogue of Eq. (1) for polarization, we define a tensor $\Delta_{ab}(s) \equiv \mathcal{P}_{ab}[x(s)] - \mathcal{P}_{||ab}(s)$ along the geodesic $x^a(s)$ connecting $\hat{\mathbf{n}}$ and its image. Here, s is distance along the geodesic measured from $\hat{\mathbf{n}}$, so $s = s_* \equiv |\alpha^a(\hat{\mathbf{n}})|$ at the end point, and $\mathcal{P}_{||ab}(s)$ is the tensor obtained by parallel transporting $\mathcal{P}_{ab}[x(s_*)]$ back along the geodesic. With these definitions, $\Delta_{ab}(s_*) = 0$ and the lensed polarization tensor at $\hat{\mathbf{n}}$ is $\tilde{\mathcal{P}}_{ab}(\hat{\mathbf{n}}) =$

*Electronic address: A.D.Challinor@mrao.cam.ac.uk

†Electronic address: gchon@mrao.cam.ac.uk

$\mathcal{P}_{ab}(\hat{\mathbf{n}}) - \Delta_{ab}(0)$. We now compute $\Delta_{ab}(s_*)$ by Taylor expanding its components Δ_{ij} in an arbitrary basis about $s = 0$. Setting the expansion to zero and using the geodesic character of $x^a(s)$, to second order we find

$$\Delta_{ij}(0) = -\alpha^k \partial_k \Delta_{ij} + \frac{1}{2} \alpha^l \alpha^m (\Gamma_{lm}^k \partial_k \Delta_{ij} - \partial_l \partial_m \Delta_{ij}), \quad (2)$$

where all terms on the right are evaluated at $s = 0$. Replacing the partial derivatives with covariant derivatives and terms involving the connection Γ_{lm}^k , and solving iteratively for $\Delta_{ab}(0)$, we find

$$\Delta_{ab}(0) = -\alpha^c \nabla_c \Delta_{ab} - \frac{1}{2} \alpha^c \alpha^d \nabla_c \nabla_d \Delta_{ab} + \dots \quad (3)$$

Finally, we use the fact that $\mathcal{P}_{\parallel ab}(s)$ is parallel transported along $x^a(s)$ to obtain our required result for the lensed polarization at $\hat{\mathbf{n}}$:

$$\tilde{\mathcal{P}}_{ab}(\hat{\mathbf{n}}) = \mathcal{P}_{ab}(\hat{\mathbf{n}}) + \alpha^c \nabla_c \mathcal{P}_{ab}(\hat{\mathbf{n}}) + \frac{1}{2} \alpha^c \alpha^d \nabla_c \nabla_d \mathcal{P}_{ab}(\hat{\mathbf{n}}) + \dots \quad (4)$$

This expansion is the obvious generalisation of Eq. (1) to the covariant transport of polarization.

The expansion of the symmetric, trace-free polarization tensor in the spin-weight formalism [3] (our conventions follow Ref. [4]) is

$$\mathcal{P}^{ab} = 2^{-2} {}_{-2}P e_+^a e_+^b + 2^{-2} {}_2P e_-^a e_-^b, \quad (5)$$

where it is convenient to choose the complex null vectors $e_\pm^a = (\partial_\theta)^a \pm i \csc \theta (\partial_\phi)^a$. With this choice, the spin ± 2 polarization ${}_{\pm 2}P = Q \mp iU$, where the Stokes parameters Q and U are measured on the $\{(\partial_\theta)^a, -\csc \theta (\partial_\phi)^a\}$ basis. With a little effort, we can establish the spin-weight version of Eq. (4):

$$\begin{aligned} {}_{-2}\tilde{P} &= {}_{-2}P - \frac{1}{2}({}_1\alpha \bar{\partial} + {}_{-1}\alpha \bar{\partial}) {}_{-2}P \\ &\quad + \frac{1}{8}({}_1\alpha {}_1\alpha \bar{\partial} \bar{\partial} + {}_1\alpha {}_{-1}\alpha \bar{\partial} \bar{\partial} \\ &\quad + {}_{-1}\alpha {}_1\alpha \bar{\partial} \bar{\partial} + {}_{-1}\alpha {}_{-1}\alpha \bar{\partial} \bar{\partial}) {}_{-2}P + \dots \end{aligned} \quad (6)$$

where ${}_{\pm 1}\alpha$ are the spin-weight components of α^a : $\alpha^a = 2^{-1} {}_1\alpha e_-^a + 2^{-1} {}_{-1}\alpha e_+^a$. In terms of the raising and lowering operators [3, 4], $\bar{\partial}$ and $\bar{\partial}$, and the lensing potential $\psi(\hat{\mathbf{n}})$, we have ${}_1\alpha = -\bar{\partial}\psi$ and ${}_{-1}\alpha = {}_1\alpha^* = -\bar{\partial}\psi$. Equation (6), which holds quite generally for the spin-weight components ${}_{\pm s}\eta$ of any symmetric, trace-free tensor $\eta_{a_1\dots a_s}$, is the correct covariant generalisation of the flat-sky result given as Eq. (43) in Ref. [1]. The difference between Eq. (6) and a simple Taylor expansion of ${}_{-2}P$, i.e. ${}_{-2}\tilde{P} = {}_{-2}P + \alpha^a \nabla_a {}_{-2}P + \dots$, amounts to a factor $e^{-2i\chi}$. This factor describes the combined effect of the rotations at $\hat{\mathbf{n}}$ and its image that are required to align one of the polarization basis vectors with the local tangent to the geodesic. The angle χ has the expansion

$$\begin{aligned} \chi &= \frac{1}{2} i e_-^a \alpha^b \nabla_b e_{+a} - \frac{1}{8} i (e_-^a \alpha^b \nabla_b e_{+a})^2 \\ &\quad + \frac{1}{4} i e_-^a \alpha^b \alpha^c \nabla_b \nabla_c e_{+a} + \dots \end{aligned} \quad (7)$$

Evaluating the covariant derivatives gives χ in terms of the (polar) coordinate components of α^a :

$$\chi = \alpha^\phi \left[\cos \theta - \frac{1}{2} \csc \theta (1 + \cos^2 \theta) \alpha^\theta + \dots \right]. \quad (8)$$

When displacing along lines of constant ϕ the rotation angle vanishes, as expected.

Having established the appropriate form of the polarization mapping describing weak lensing on the sphere, it is straightforward to compute the power spectra of the lensed fields. One can either work directly with the polarization tensor, using Eq. (5) and an expansion in the tensor harmonics [5], or proceed in the spin-weight formalism using Eq. (6) and an expansion in spin-weight harmonics [6]. Here we shall follow the latter route and sketch a rigorous derivation of the polarization power spectra. Extracting the spin- ± 2 components, ${}_{\pm 2}\tilde{P}_{lm}$, of the lensed polarization using Eq. (6) we find

$$\begin{aligned} {}_{\pm 2}\tilde{P}_{lm} &= {}_{\pm 2}P_{lm} + \psi_{(lm)_1} {}_{\pm 2}P_{(lm)_2} + {}_{\pm 2}I_{ll_1l_2}^{mm_1m_2} \\ &\quad + \frac{1}{2} \psi_{(lm)_1} \psi_{(lm)_2}^* {}_{\pm 2}P_{(lm)_3} + {}_{\pm 2}J_{ll_1l_2l_3}^{mm_1m_2m_3}, \end{aligned} \quad (9)$$

where summation should be understood over repeated multipole indices (l and m). Here ${}_{\pm 2}P_{lm}$ are the spin-weight multipoles of the unlensed polarization and ψ_{lm} are the usual spherical multipoles of the scalar lensing potential. The overlap integrals are

$$\begin{aligned} {}_{\pm 2}I_{ll_1l_2}^{mm_1m_2} &\equiv \int d\hat{\mathbf{n}} {}_{\pm 2}Y_{lm}^* \frac{1}{2} (\bar{\partial} Y_{(lm)_1} \bar{\partial} + \bar{\partial} Y_{(lm)_1} \bar{\partial}) \\ &\quad \times {}_{\pm 2}Y_{(lm)_2}, \end{aligned} \quad (10)$$

$$\begin{aligned} {}_{\pm 2}J_{ll_1l_2l_3}^{mm_1m_2m_3} &\equiv \int d\hat{\mathbf{n}} {}_{\pm 2}Y_{lm}^* \frac{1}{4} [\bar{\partial} Y_{(lm)_1} \bar{\partial} Y_{(lm)_2}^* \bar{\partial} \bar{\partial} \\ &\quad + \bar{\partial} Y_{(lm)_1} \bar{\partial} Y_{(lm)_2}^* \bar{\partial} \bar{\partial} \\ &\quad + \bar{\partial} Y_{(lm)_1} \bar{\partial} Y_{(lm)_2}^* (\bar{\partial} \bar{\partial} + \bar{\partial} \bar{\partial})] \\ &\quad \times {}_{\pm 2}Y_{(lm)_3}, \end{aligned} \quad (11)$$

which should be compared with Eq. (65) in Ref. [1]¹. The integral defining ${}_{\pm 2}I_{ll_1l_2}^{mm_1m_2}$ can be evaluated by integrating by parts using the spin-weight integral theorems in the appendices of Ref. [4], and noting that the spin-weight harmonics ${}_sY_{lm}$ are eigenfunctions of $\bar{\partial} \bar{\partial}$ with eigenvalue $s(s-1) - l(l-1)$. [The identity $(\bar{\partial} \bar{\partial} - \bar{\partial} \bar{\partial}) {}_s\eta = 2 {}_s\eta$ for spin- s ${}_s\eta$ is also useful.] The integral for ${}_{\pm 2}I_{ll_1l_2}^{mm_1m_2}$ then reduces to the right-hand side of Eq. (72) in Ref. [1], which can be evaluated in terms of 3- j symbols.

Breaking the spin-weight multipoles into parity eigenstates, ${}_{\pm 2}P_{lm} = E_{lm} \pm iB_{lm}$, where E_{lm} are the multipoles of electric polarization and B_{lm} of magnetic, and assuming no correlations between polarization and the lensing potential, the overlap integrals ${}_{\pm 2}I$ enter the correlators such as $\langle \tilde{E}_{lm} \tilde{E}_{(lm)'}^* \rangle$ in the form $\sum_{m_1m_2} {}_2I_{ll_1l_2}^{mm_1m_2} {}_2I_{l'l_1l_2}^{m'm_1m_2*}$. The orthogonality relations

¹ There is a typo in the second line in Eq. (65) of Ref. [1]: one of the derivatives acting on the spin-0 harmonics should be ∇_j rather than ∇_i .

for the $3\text{-}j$ symbols ensure that the summation vanishes unless $l = l'$ and $m = m'$ as required to maintain statistical isotropy of the lensed fields. The overlap integrals ${}_{\pm 2}J$ enter the correlators in the form $\sum_{m_1} {}_{\pm 2}J_{l_1 l_1'}^{m m_1 m_1'}$ (and its complex conjugate). Setting $m_1 = m_2$ in Eq. (11), and summing, isolates terms

$$\sum_{m_1} \bar{\partial} Y_{(lm)_1} \bar{\partial} Y_{(lm)_1}^* = l_1(l_1 + 1) \frac{2l_1 + 1}{4\pi}, \quad (12)$$

and $\sum_{m_1} \bar{\partial} Y_{(lm)_1} \bar{\partial} Y_{(lm)_1}^* = 0$, where we have used the addition theorem for spin-weight harmonics. It follows that only the term involving the operator $\bar{\partial} \bar{\partial} + \bar{\partial} \bar{\partial}$ in ${}_{\pm 2}J$ survives the summation. Finally, making use of

$$\frac{1}{2}(\bar{\partial} \bar{\partial} + \bar{\partial} \bar{\partial})_s Y_{lm} = [s^2 - l(l + 1)]_s Y_{lm}, \quad (13)$$

we find

$$\sum_{m_1} {}_{\pm 2}J_{l_1 l_1'}^{m m_1 m_1'} = -\frac{1}{2}[l(l+1)-4]l_1(l_1+1)\frac{2l_1+1}{4\pi}\delta_{ll'}\delta_{mm'}, \quad (14)$$

as required to maintain statistical isotropy. Putting these results together gives the final expressions for the lensed power spectra which agree with those given by Hu as Eq. (76) in Ref. [1]. Our result for the cross power spectrum between E and Θ also agrees with Ref. [1].

We close by discussing the relation of our derivation to that in Ref. [1]. The critical point to note is that if

the covariant derivatives there, acting on a spin-weight s (≥ 0) quantity ${}_s\eta$, are interpreted as encoding the operation $2^{-s}e_+^{A_s}\nabla_b({}_s\eta e_{-A_s})$, where $e_{\pm}^{A_s}$ denotes the irreducible tensor product $e_{\pm}^{a_1}\dots e_{\pm}^{a_s}$, then the lens mapping implied by Eq. (64) of Ref. [1] reduces to that given here. (The result that $e_+^{A_s}e_{-B_s}$ is covariantly constant proves useful.) Furthermore, it can be shown that the integration by parts used to simplify the overlap integrals ${}_{\pm 2}J$ remains consistent under this interpretation of the covariant derivative acting on spin-weight quantities. Unfortunately, this compressed notation appears to have been adopted in Ref. [1] without comment. [It is hinted at in Eq. (55) there, and in the comment above Eq. (71).] Since the raising and lowering operators of the spin-weight formalism naturally encode the required derivative operations, there is no need to overload the standard notation of differential geometry. Furthermore, the overly-compact notation of Ref. [1] hides the covariant nature of the lens re-mapping of polarization on the sphere.

Acknowledgments

AC acknowledges a PPARC Postdoctoral Fellowship. We thank Wayne Hu for useful comments.

-
- [1] W. Hu, Phys. Rev. D **62**, 043007 (2000).
 - [2] N. Kaiser, Astrophys. J. **498**, 26 (1998).
 - [3] J. N. Goldberg, A. J. MacFarlane, E. T. Newman, F. Rohrlich, and E. C. G. Sudarshan, J. Math. Phys. **8**, 2155 (1967).
 - [4] A. Lewis, A. Challinor, and N. Turok, Phys. Rev. D **65**,

- 023505 (2002).
- [5] M. Kamionkowski, A. Kosowsky, and A. Stebbins, Phys. Rev. Lett. **78**, 2058 (1997).
- [6] U. Seljak and M. Zaldarriaga, Phys. Rev. Lett. **78**, 2054 (1997).